

## 4B PREP INSTRUCTIONS

8 days before procedure	7 days before procedure	3 days before procedure	2 days before procedure
Date	Date		
<input type="checkbox"/> Stop any weekly injectable weight loss medications (ex- Ozempic, Wegovy, Zepbound) Arrange for a friend, family member or medical transport to drive you home after the colonoscopy.	Start low residue diet Buy preparation ingredients <input type="checkbox"/> One gallon of GoLyteLy or NuLyteLy (4 liters) Rx sent to your pharmacy <input type="checkbox"/> 4 Simethicone (Gas-X) Tablets (80mg each) Rx sent to your pharmacy <input type="checkbox"/> 6 packets of Miralax (17g per packet), over the counter	Continue low residue diet After lunch Mix 2 packets of miralax (or measure out 34g) in any 8 to 16 oz of water, Gatorade or plain juice, then drink.	Continue low residue diet After lunch Mix 2 packets of miralax (or measure out 34g) in any 8 to 16 oz of water, Gatorade or plain juice, then drink.
1 day before procedure	Day of Procedure	What to Expect with your Prep	
Date  No Marijuana or Alcohol  Breakfast- Start clear liquid diet and continue until procedure  Stay hydrated with at least 12 glasses of clear liquids throughout the day (in addition to what you drink with your bowel prep medication)  12pm Mix 2 packets of MiraLAX (or measure out 34g), in water, Gatorade or juice and drink.  5pm <input type="checkbox"/> Take 2 Simethicone tablets (Gas-X) <input type="checkbox"/> Separate GoLyteLy/NuLyteLy solution in half. <input type="checkbox"/> Begin drinking one half of GoLyteLy/NuLyteLy (shake well). Drink a large glass of the prep (8oz) every 15 minutes. It is best to drink the whole glass rapidly rather than sipping slowly. <input type="checkbox"/> Once half the gallon of GoLyteLy/NuLyteLy is complete, take the last 2 Simethicone tablets	Date  No tobacco or marijuana use  5am GoLyteLy/NuLyteLy (shake well). Drink a large glass of the prep (8oz) every 15 minutes. It is best to drink the whole glass rapidly rather than sipping slowly. You must finish this no later than 4 hours prior to your arrival time. Do not eat! No gum, mints or cough drops!  Stop Clear Liquids at _____  Arrival time _____  After your procedure You should not drive a car, operate machinery, or make any legal decisions until the day after your procedure.  You must have an adult accompany you home the day of your procedure. The staff must be able to contact whoever will accompany you.	You will develop significant diarrhea after drinking the preparation. Plan to be near a bathroom. This is normal as it means the medication is working to clear stool from your colon.  Most people feel mild bloating and may experience abdominal cramps. This is normal. Drinking the prep medication more slowly and over a longer period of time can help alleviate these symptoms.  A successful colon prep will cause you to have clear yellow ("tea colored") liquid stools.  Please finish your preparation regardless of your stool color/form.	

**FAILURE TO FOLLOW THESE INSTRUCTIONS MAY RESULT IN CANCELLATION OF YOUR PROCEDURE**